

**THE QUANTUM SPACES OF CERTAIN GRADED ALGEBRAS
RELATED TO $\mathfrak{sl}(2, \mathbb{k})$**

RICHARD G. CHANDLER*

richard.chandler@untdallas.edu

www.untdallas.edu/faculty/chandler

Department of Mathematics and Information Sciences,

University of North Texas at Dallas,

Dallas, TX 75241

and

MICHAELA VANCLIFF*

vancliff@uta.edu

www.uta.edu/math/vancliff

Department of Mathematics,

University of Texas at Arlington,

Arlington, TX 76019

ABSTRACT. Inspired by the work of Le Bruyn and Smith in [12] and the work of Shelton and Vancliff in [17], we analyze certain graded algebras related to the Lie algebra $\mathfrak{sl}(2, \mathbb{k})$ using geometric techniques in the spirit of Artin, Tate and Van den Bergh. In particular, we discuss the point schemes and line schemes of certain quadratic quantum \mathbb{P}^3 s associated to the Lie superalgebra $\mathfrak{sl}(1|1)$, to a quantized enveloping algebra, $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$, of $\mathfrak{sl}(2, \mathbb{k})$, and to a color Lie algebra $\mathfrak{sl}_k(2, \mathbb{k})$, respectively. The geometry we consider identifies the existence of a central element in the universal enveloping algebra of $\mathfrak{sl}(1|1)$ and in $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$.

INTRODUCTION

A regular algebra of global dimension four is often viewed as a noncommutative analogue of a polynomial ring on four variables. In the language of [4], such an algebra is frequently called a quantum \mathbb{P}^3 .

In [12], the point modules and line modules, called the quantum space, of the homogenization $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$ of the universal enveloping algebra of the Lie algebra $\mathfrak{sl}(2, \mathbb{k})$ were determined.

2010 *Mathematics Subject Classification.* 14A22, 16S37, 17B35, 17B37, 17B75.

Key words and phrases. line scheme, point scheme, Lie algebra, superalgebra, regular algebra, Plücker coordinates.

*This work was supported in part by NSF grants DMS-0900239 and DMS-1302050.

Therein, it was shown that the scheme of points in \mathbb{P}^3 that correspond to the point modules of $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$ is the union of a point, a plane, and a conic embedded in the plane, whereas the lines in \mathbb{P}^3 that correspond to the line modules of $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$ are the lines lying on a certain pencil of quadrics. The quantum space of $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$ identifies certain features of $\mathfrak{sl}(2, \mathbb{k})$, including the existence of a Casimir element in the universal enveloping algebra.

The definition of line scheme is given in [16]. In [17], a method was given for computing the line scheme of any quadratic algebra on four generators that is a domain and has a Hilbert series the same as that of the polynomial ring on four variables. Therein, it was shown that the line scheme of $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$ is neither irreducible nor reduced and has two two-dimensional components. In contrast, the computation of the line modules in [12] was done without the use of the line scheme.

Following the work in [12, 17], we analyze a quantum space associated to the Lie superalgebra $\mathfrak{sl}(1|1)$ (cf. [6]), to a quantized enveloping algebra, $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$, of $\mathfrak{sl}(2, \mathbb{k})$ (cf. [18]), and to the color Lie algebra $\mathfrak{sl}_k(2, \mathbb{k})$ defined in [5]. As in [12], the quantum space we associate to each of the first two algebras identifies some distinguished elements in the associated graded algebra.

The article is outlined as follows. In Section 1, we will discuss some background material relevant to later sections. In Section 2, we analyze $\mathfrak{sl}(1|1)$ by first computing the point scheme and line variety of an associated quadratic quantum \mathbb{P}^3 , denoted $\mathcal{H}(\mathfrak{sl}(1|1))$. We then identify the lines in \mathbb{P}^3 that correspond to the line modules of $\mathcal{H}(\mathfrak{sl}(1|1))$ and then identify $\mathcal{H}(\mathfrak{sl}(1|1))$ as a twist by an automorphism of a member of the family of coordinate rings of quantum 2×2 matrices [8]; this aids in recognizing a certain element of $\mathcal{H}(\mathfrak{sl}(1|1))$ as a distinguished central element. In Section 3, we analyze $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$ in a similar manner using an associated quadratic quantum \mathbb{P}^3 , denoted $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$. The geometry identifies an element in $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ related to the quantum Casimir element in $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$. In Section 4, we examine $\mathfrak{sl}_k(2, \mathbb{k})$ via an associated quadratic quantum \mathbb{P}^3 , denoted $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$. The geometry associated to $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ suggests the lack of a distinguished central element akin to those found in the analysis of $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$, $\mathcal{H}(\mathfrak{sl}(1|1))$, and $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$. Section 5 is an Appendix which contains relevant polynomials and matrices used in the computations of the various point schemes and line varieties in the article.

1. PRELIMINARIES

Throughout this article, \mathbb{N} denotes the set of positive integers, \mathbb{k} denotes an algebraically closed field of characteristic zero, $\mathbb{k}^\times = \mathbb{k} \setminus \{0\}$, $\mathcal{V}(f_1, \dots, f_m)$ denotes the zero locus of the polynomials f_1, \dots, f_m , and e_1, \dots, e_4 denote the points $e_1 = (1, 0, 0, 0)$, $e_2 = (0, 1, 0, 0)$, $e_3 = (0, 0, 1, 0)$ and $e_4 = (0, 0, 0, 1)$ in \mathbb{P}^3 .

Given a quadratic regular algebra, A , on degree-one generators x_1, x_2, x_3, x_4 , we may follow [1] in order to compute the point scheme of A . We write the defining relations of A in the form

$Mx = 0$, where M is 6×4 matrix and x is a column vector given by $x^T = (x_1, x_2, x_3, x_4)$. The point scheme of A can be identified with the zero locus in $\mathbb{P}(A_1^*) \cong \mathbb{P}^3$ of the 4×4 minors of M .

Our computation of the line scheme as a subscheme of \mathbb{P}^5 will make use of the Plücker coordinates, $M_{12}, M_{13}, M_{14}, M_{23}, M_{24}, M_{34}$ on \mathbb{P}^5 . We first recall how these coordinates relate to lines in \mathbb{P}^3 ; details may be found in [7, §8.6].

Any line ℓ in \mathbb{P}^3 is uniquely determined by any two distinct points $(a_1, a_2, a_3, a_4), (b_1, b_2, b_3, b_4)$ belonging to ℓ ; so ℓ may be represented by a 2×4 matrix

$$\begin{bmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \end{bmatrix}$$

that has rank two; in particular, the points on ℓ are represented in homogeneous coordinates by linear combinations of the rows of this matrix.

The Plücker coordinate M_{ij} is evaluated on this matrix as the minor $a_i b_j - a_j b_i$, for all $i < j$, and the Plücker polynomial $P = M_{12}M_{34} - M_{13}M_{24} + M_{14}M_{23}$ vanishes on this matrix. Moreover, $\mathcal{V}(P)$ is the subscheme of \mathbb{P}^5 that parametrizes all the lines in \mathbb{P}^3 .

2. THE LIE SUPERALGEBRA $\mathfrak{sl}(1|1)$

2.1. The \mathbb{k} -Algebra $\mathcal{H}(\mathfrak{sl}(1|1))$.

Let $\mathfrak{sl}(1|1) = \mathbb{k}e \oplus \mathbb{k}f \oplus \mathbb{k}h$ denote the Lie superalgebra (cf. [6]) with bracket defined by

$$[e, f] = h, \quad [h, e] = [h, f] = [e, e] = [f, f] = [h, h] = 0.$$

The universal enveloping algebra (cf. [6]) of $\mathfrak{sl}(1|1)$ is

$$\mathcal{U}(\mathfrak{sl}(1|1)) = \frac{\mathbb{k}\langle e, f, h \rangle}{\langle ef + fe - h, he - eh, hf - fh, e^2, f^2 \rangle}.$$

In order to obtain a quadratic quantum \mathbb{P}^3 that maps onto $\mathcal{U}(\mathfrak{sl}(1|1))$, we define the following algebra. In Theorem 2.2, we establish that the algebra is indeed a quadratic quantum \mathbb{P}^3 .

Definition 2.1. The quadratic \mathbb{k} -algebra, $\mathcal{H}(\mathfrak{sl}(1|1))$, on degree-one generators e, f, h, t is

$$\mathcal{H}(\mathfrak{sl}(1|1)) = \frac{\mathbb{k}\langle e, f, h, t \rangle}{\langle ef + fe - ht, he - eh, hf - fh, et - te, ft - tf, ht - th \rangle}.$$

Note that $\mathcal{U}(\mathfrak{sl}(1|1)) \cong \mathcal{H}(\mathfrak{sl}(1|1)) / \langle t - 1, e^2, f^2 \rangle$.

Theorem 2.2. *The \mathbb{k} -algebra $\mathcal{H}(\mathfrak{sl}(1|1))$ is a quadratic quantum \mathbb{P}^3 .*

Proof. Let $A = \mathbb{k}[e, h, t]$. The algebra $\mathcal{H}(\mathfrak{sl}(1|1))$ is isomorphic to the Ore extension (cf. [9]) $A[f; \sigma, \delta]$, where σ and δ are an automorphism and a σ -derivation, respectively, of A defined by

$$\sigma(e) = -e, \quad \sigma(h) = h, \quad \sigma(t) = t, \quad \delta(e) = ht, \quad \delta(h) = 0, \quad \delta(t) = 0.$$

Hence, by [15], $\mathcal{H}(\mathfrak{sl}(1|1))$ is Auslander-regular and satisfies the Cohen-Macaulay property. By the definition of the Cohen-Macaulay property [14, Definition 5.8], $\mathcal{H}(\mathfrak{sl}(1|1))$ has polynomial growth and is, hence, AS-regular by [14]. By [14, Theorem 4.8], $\mathcal{H}(\mathfrak{sl}(1|1))$ is a domain and, thus, t is a central regular element of $\mathcal{H}(\mathfrak{sl}(1|1))$. It follows that $\mathcal{H}(\mathfrak{sl}(1|1))$ is a central regular extension of $\mathbb{k}[e, f, h]$, in the language of [13], and so is AS-regular of global dimension four (cf. [13, Theorem 2.6, Corollary 2.7] and the paragraph after [13, Definition 3.1.1]). Hence, $\mathcal{H}(\mathfrak{sl}(1|1))$ is a quadratic quantum \mathbb{P}^3 . \blacksquare

Note that there exists an automorphism φ on $\mathcal{H}(\mathfrak{sl}(1|1))$ defined by $\varphi : e \leftrightarrow f$.

2.2. The Quantum Space of $\mathcal{H}(\mathfrak{sl}(1|1))$.

Denote $x_1 = e$, $x_2 = f$, $x_3 = h$ and $x_4 = t$.

Theorem 2.3. *The point scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$ is isomorphic to the subscheme*

$$\mathfrak{p} = \mathcal{V}(x_3x_4 - 2x_1x_2) \cup \mathcal{V}(x_3, x_4)$$

of \mathbb{P}^3 ; that is, \mathfrak{p} is the union of a nonsingular quadric and a line in \mathbb{P}^3 that meets the quadric at two distinct points.

Proof. The polynomials that define \mathfrak{p} are listed in Appendix 5.1. The zero locus of these polynomials is $\mathcal{V}(x_3x_4 - 2x_1x_2) \cup \mathcal{V}(x_3, x_4)$.

The Jacobian matrix, J , of \mathfrak{p} is given in Appendix 5.2. If p is a multiple point contained in a d -dimensional, irreducible, component of \mathfrak{p} , then the $(3-d) \times (3-d)$ minors of $J|_p$ vanish, and conversely [10]. The only such points in \mathfrak{p} are $e_1, e_2 \in \mathcal{V}(x_3x_4 - 2x_1x_2) \cap \mathcal{V}(x_3, x_4)$.

The coordinate ring of \mathfrak{p} is $\mathbb{k}[x_1, x_2, x_3, x_4]/I$, where I is the ideal generated by the polynomials in Appendix 5.1. The line $L = \mathcal{V}(x_2 - x_3, x_4)$ is a hyperplane of complementary dimension to $\mathcal{V}(x_3x_4 - 2x_1x_2)$ in \mathbb{P}^3 and the points of intersection of L and \mathfrak{p} are e_1 and $(0, 1, 1, 0)$. Localizing the coordinate ring of $\mathfrak{p} \cap L$ around e_1 yields a ring isomorphic to $\mathbb{k}[x]/\langle x^2 \rangle$ and so e_1 has multiplicity two. This implies that e_1 is a multiple point of \mathfrak{p} only as a consequence of being an intersection point of the two irreducible components of \mathfrak{p} . Applying the automorphism φ allows us to conclude that the same applies to e_2 . Hence, \mathfrak{p} is as proposed. \blacksquare

Corollary 2.4. *Let $A = \mathcal{H}(\mathfrak{sl}(1|1))$.*

- (i) *The points in $\mathbb{P}(A_1^*) \times \mathbb{P}(A_1^*)$ on which the defining relations of $\mathcal{H}(\mathfrak{sl}(1|1))$ vanish are of the form (p, p) , if $p \in \mathcal{V}(x_3x_4 - 2x_1x_2)$ and are of the form $((\alpha_1, \alpha_2, 0, 0), (\alpha_1, -\alpha_2, 0, 0))$, if $(\alpha_1, \alpha_2, 0, 0) \in \mathcal{V}(x_3, x_4)$.*
- (ii) *There exists an automorphism $\sigma : \mathfrak{p} \rightarrow \mathfrak{p}$ which, on closed points, is defined by*

$$\sigma(p) = \sigma(p_1, p_2, p_3, p_4) = \begin{cases} (p_1, p_2, p_3, p_4), & p \in \mathcal{V}(x_3x_4 - 2x_1x_2) \\ (p_1, -p_2, 0, 0), & p \in \mathcal{V}(x_3, x_4). \end{cases}$$

Proof. Part (i) is shown by computation. The existence of the map in (ii) follows from (i) and [13]. \blacksquare

Theorem 2.5. *The line scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$ has dimension three and its reduced variety, \mathfrak{L} , is the union of the irreducible components:*

- (I) $\mathfrak{L}_1 = \mathcal{V}(M_{34}, M_{13}M_{24} - M_{14}M_{23})$,
- (II) $\mathfrak{L}_2 = \mathcal{V}(M_{14}, M_{23}, M_{34}^2 + 2M_{13}M_{24}, 2M_{12} + M_{34})$,
- (III) $\mathfrak{L}_3 = \mathcal{V}(M_{13}, M_{24}, M_{34}^2 + 2M_{14}M_{23}, 2M_{12} - M_{34})$.

Proof. For details on the construction of the line scheme from the defining relations of a quadratic quantum \mathbb{P}^3 , the reader is referred to [4, 17].

The polynomials that define the line scheme are given in Appendix 5.3. The reduced variety given by those polynomials is \mathfrak{L} . For details of the computation, the reader is referred to [3]. \blacksquare

Corollary 2.6. *The lines in \mathbb{P}^3 that are parametrized by the closed points of the line scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$ are:*

- (i) *the lines in \mathbb{P}^3 that intersect the line $\mathcal{V}(x_3, x_4)$, and*
- (ii) *the lines in \mathbb{P}^3 that lie on the quadric $\mathcal{V}(2x_1x_2 - x_3x_4)$.*

Proof. Let $(a_1, a_2, a_3, a_4), (b_1, b_2, b_3, b_4) \in \mathbb{P}^3$ be distinct points and let

$$\ell = \begin{bmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \end{bmatrix}$$

represent the line between them.

- (i) If ℓ is given by a point of \mathfrak{L}_1 , then it follows that ℓ is represented by a rank-two matrix of the form

$$\begin{bmatrix} c_1 & c_2 & c_3 & c_4 \\ c_5 & c_6 & 0 & 0 \end{bmatrix},$$

where $c_1, \dots, c_6 \in \mathbb{k}$. Hence, ℓ intersects $\mathcal{V}(x_3, x_4)$ if and only if ℓ is given by \mathfrak{L}_1 .

- (ii) If ℓ is given by \mathfrak{L}_2 , then it follows that ℓ may be represented by

$$\begin{bmatrix} a_1 & 0 & 0 & a_4 \\ 0 & a_2 & a_3 & 0 \end{bmatrix},$$

where $(a_1, a_4), (a_2, a_3) \in \mathbb{P}^1$ and $2a_1a_2 - a_3a_4 = 0$. Both $(a_1, 0, 0, a_4)$ and $(0, a_2, a_3, 0)$ lie on the quadric $\mathcal{V}(2x_1x_2 - x_3x_4)$; in fact, for any point $p = (a_1, \delta a_2, \delta a_3, a_4) \in \ell$, where $\delta \in \mathbb{P}^1$, p belongs to $\mathcal{V}(2x_1x_2 - x_3x_4)$. Thus, ℓ belongs to one of the rulings of the quadric $\mathcal{V}(2x_1x_2 - x_3x_4)$. In particular, $\ell \in \{\mathcal{V}(\mu x_1 - x_4, 2x_2 - \mu x_3) : \mu \in \mathbb{P}^1\}$.

The component \mathfrak{L}_3 may be obtained from \mathfrak{L}_2 by applying φ ; thus, ℓ is given by \mathfrak{L}_3 and only if ℓ belongs to the ruling $\{\mathcal{V}(\mu x_1 - x_3, 2x_2 - \mu x_4) : \mu \in \mathbb{P}^1\}$ of $\mathcal{V}(2x_1x_2 - x_3x_4)$.

Since every line on $\mathcal{V}(2x_1x_2 - x_3x_4)$ belongs to exactly one of the rulings above, it follows that the lines on $\mathcal{V}(2x_1x_2 - x_3x_4)$ are in one-to-one correspondence with the points in $\mathfrak{L}_2 \cup \mathfrak{L}_3$. \blacksquare

2.3. Twisting $\mathcal{O}_q(\mathbb{M}_2)$.

Given the results of Section 2.2, the quantum space of $\mathcal{H}(\mathfrak{sl}(1|1))$ is isomorphic to the quantum space of the coordinate ring, $\mathcal{O}_q(\mathbb{M}_2)$, of quantum 2×2 matrices (see Definition 2.7). This fact suggests that $\mathcal{H}(\mathfrak{sl}(1|1))$ could be isomorphic to a twist by an automorphism of $\mathcal{O}_q(\mathbb{M}_2)$, for some $q \in \mathbb{k}^\times$.

Definition 2.7. [8] The coordinate ring of quantum 2×2 matrices is the \mathbb{k} -algebra

$$\mathcal{O}_q(\mathbb{M}_2) = \frac{\mathbb{k}\langle a, b, c, d \rangle}{\langle ab - qba, cd - qdc, ac - qca, bd - qdb, bc - cb, ad - da - (q - q^{-1})bc \rangle},$$

where $q \in \mathbb{k}^\times, q^2 \neq 1$.

Lemma 2.8. [19]

- (i) *The point scheme of $\mathcal{O}_q(\mathbb{M}_2)$ is $\mathcal{V}(ad - bc) \cup \mathcal{V}(b, c)$.*
- (ii) *The closed points of the line scheme of $\mathcal{O}_q(\mathbb{M}_2)$ parametrize all the lines in \mathbb{P}^3 that belong to $\mathcal{V}(ad - bc)$ and all lines that intersect $\mathcal{V}(b, c)$.*

Theorem 2.9. *The algebra $\mathcal{H}(\mathfrak{sl}(1|1))$ is isomorphic to a twist by an automorphism (in the sense of [2]) of the algebra $\mathcal{O}_i(\mathbb{M}_2)$, where $i^2 = -1$.*

Proof. Let $A = \mathcal{O}_i(\mathbb{M}_2)$ and $\tau : A \rightarrow A$ be the automorphism defined by

$$\tau(a) = ia, \quad \tau(b) = b, \quad \tau(c) = c, \quad \tau(d) = -id.$$

Also, let A^τ be the algebra obtained by twisting A by τ , with multiplication, \star , defined by $\bar{x} \star \bar{y} = x\tau(y)$, for all $\bar{x}, \bar{y} \in A_1^\tau$, where \bar{x}, \bar{y} are the elements of A_1^τ corresponding to x and y in A_1 . It follows that:

$$A^\tau \cong \frac{\mathbb{k}\langle a, b, c, d \rangle}{\langle ab - ba, ac - ca, bc - cb, bd - db, cd - dc, ad + da - 2bc \rangle}.$$

This algebra is isomorphic to $\mathcal{H}(\mathfrak{sl}(1|1))$ under the isomorphism $\psi : A^\tau \rightarrow \mathcal{H}(\mathfrak{sl}(1|1))$ defined by

$$\psi(a) = e, \quad \psi(b) = \frac{h}{\sqrt{2}}, \quad \psi(c) = \frac{t}{\sqrt{2}}, \quad \psi(d) = f. \quad \blacksquare$$

This result leads to the identification of a distinguished supercommuting element of $\mathcal{H}(\mathfrak{sl}(1|1))$ (and therefore of $\mathcal{U}(\mathfrak{sl}(1|1))$). The element $ad - ibc \in \mathcal{O}_i(\mathbb{M}_2)$, called the quantum determinant, is central in $\mathcal{O}_i(\mathbb{M}_2)$; its image in $\mathcal{H}(\mathfrak{sl}(1|1))$ belongs to $\mathbb{k}^\times(2ef - ht)$, which corresponds to the

quadric $\mathcal{V}(2ef - ht)$ in the point scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$ since $2ef - ht$ vanishes on the graph of $\sigma|_{\mathcal{V}(2ef - ht)}$.

Regrading $\mathcal{H}(\mathfrak{sl}(1|1))$ by taking e and f to be odd and h and t to be even, $2ef - ht$ supercommutes in the algebra. The image of this element in $\mathcal{U}(\mathfrak{sl}(1|1))$ is $2ef - h = ef - fe$ which supercommutes in $\mathcal{U}(\mathfrak{sl}(1|1))$. More precisely, $\mathcal{U}(\mathfrak{sl}(1|1))$ has a PBW basis given by $\{e^i f^j h^k : i, j, k \in \mathbb{N} \cup \{0\}\}$ and can be realized as a superalgebra by taking

$$\mathcal{U}(\mathfrak{sl}(1|1))_0 = \bigoplus_{i+j+k \text{ is even}} \mathbb{k}(e^i f^j h^k), \quad \mathcal{U}(\mathfrak{sl}(1|1))_1 = \bigoplus_{i+j+k \text{ is odd}} \mathbb{k}(e^i f^j h^k).$$

If $x \in \mathcal{U}(\mathfrak{sl}(1|1))_0$, then $x(2ef - h) = (2ef - h)x$; if $x \in \mathcal{U}(\mathfrak{sl}(1|1))_1$, then $x(2ef - h) = -(2ef - h)x$.

Hence, one may view the element $2ef - h \in \mathcal{U}(\mathfrak{sl}(1|1))$ as playing the role of a Casimir element of a Lie superalgebra, which is not a well-defined concept for $\mathcal{U}(\mathfrak{sl}(1|1))$ since the Killing Form on $\mathfrak{sl}(1|1)$ is degenerate. Hence, the geometry we associated to $\mathcal{H}(\mathfrak{sl}(1|1))$ was able to identify a ‘‘generalized Casimir’’ element of $\mathcal{U}(\mathfrak{sl}(1|1))$.

3. THE QUANTIZED UNIVERSAL ENVELOPING ALGEBRA OF $\mathfrak{sl}(2, \mathbb{k})$

3.1. The \mathbb{k} -Algebra $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$.

Let $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$ denote the \mathbb{k} -algebra, known as quantized $\mathfrak{sl}(2, \mathbb{k})$, defined as

$$\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k})) = \frac{\mathbb{k}\langle E, F, K, K^{-1} \rangle}{\left\langle KE - q^2 EK, KF - q^{-2} FK, EF - FE - \frac{K^2 - K^{-2}}{q^2 - q^{-2}} \right\rangle},$$

where $q \in \mathbb{k}^\times$ and $q^4 \neq 1$ (cf. [18]).

It should be noted that this is not the current, official version of quantized $\mathfrak{sl}(2, \mathbb{k})$; that version replaces the relation $EF - FE - \frac{K^2 - K^{-2}}{q^2 - q^{-2}} = 0$ with $EF - FE - \frac{K - K^{-1}}{q - q^{-1}} = 0$ in the defining relations (cf. [11]). We thank S. P. Smith of the University of Washington for bringing to our attention that if A denotes the graded algebra defined below in Definition 3.1 and if \mathcal{O}_q denotes the current official version of quantized $\mathfrak{sl}(2, \mathbb{k})$ (cf. [11]), then the ring of degree-zero elements in $A[(KT)^{-1}]$ is isomorphic to \mathcal{O}_c , where $c^2 = q$.

Definition 3.1. We define $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ to be the quadratic \mathbb{k} -algebra on degree-one generators E, F, K, T with defining relations

$$\begin{aligned} KT &= TK, & KE &= q^2 EK, & KF &= q^{-2} FK, \\ ET &= q^2 TE, & FT &= q^{-2} TF, & EF - FE &= \frac{K^2 - T^2}{q^2 - q^{-2}}, \end{aligned}$$

where $q \in \mathbb{k}^\times$ and $q^4 \neq 1$.

Note that $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k})) \cong \mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k})) / \langle KT - 1 \rangle$. We will establish in Theorem 3.2 that $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is a quadratic quantum \mathbb{P}^3 .

Theorem 3.2. *The \mathbb{k} -algebra $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is a quadratic quantum \mathbb{P}^3 .*

Proof. Let A denote the \mathbb{k} -algebra on generators E, K, T with defining relations

$$KT = TK, \quad EK = q^{-2}KE, \quad ET = q^2TE.$$

The algebra $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is isomorphic to the Ore extension (cf. [9]) $A[F; \sigma, \delta]$, where σ and δ are an automorphism and a σ -derivation, respectively, of A defined by

$$\sigma(E) = E, \quad \sigma(K) = q^2K, \quad \sigma(T) = q^{-2}T, \quad \delta(E) = \frac{T^2 - K^2}{q^2 - q^{-2}}, \quad \delta(K) = \delta(T) = 0.$$

By [15], since A is a skew polynomial ring, $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is Auslander-regular and satisfies the Cohen-Macaulay property. By the definition of the Cohen-Macaulay property [14, Definition 5.8], $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ has polynomial growth and is, hence, AS-regular. By [14, Theorem 4.8], $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is a domain and, thus, T is a normal regular element of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$. It follows that $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is a normal regular extension of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k})) / \langle T \rangle$ (in the language of [13]), which is an Ore extension of a skew polynomial ring, so $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is AS-regular of global dimension four (cf. [13, Theorem 2.6, Corollary 2.7] and the paragraph after [13, Definition 3.1.1]). Hence, $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is a quadratic quantum \mathbb{P}^3 . ■

Note that there exists an automorphism φ on $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ defined by $\varphi : E \leftrightarrow F$ and $K \leftrightarrow T$.

3.2. The Quantum Space of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$.

Denote $x_1 = E$, $x_2 = F$, $x_3 = K$ and $x_4 = T$.

Theorem 3.3. *For every $q \in \mathbb{k}^\times$ with $q^4 \neq 1$, the point scheme, $\mathfrak{p}(q)$, of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is the union of a line, two conics and two points:*

- (i) $\mathfrak{p}_1 = \mathcal{V}(x_3, x_4)$,
- (ii) $\mathfrak{p}_2 = \mathcal{V}(x_3, q^4x_4^2 + (q^4 - 1)^2x_1x_2)$,
- (iii) $\mathfrak{p}_3 = \mathcal{V}(x_4, q^4x_3^2 + (q^4 - 1)^2x_1x_2)$,
- (iv) $\mathfrak{p}_4 = \mathcal{V}(x_1, x_2, x_3 + x_4)$,
- (v) $\mathfrak{p}_5 = \mathcal{V}(x_1, x_2, x_3 - x_4)$.

Proof. The point scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is computed in the same manner as in Theorem 2.3. The polynomials which define the scheme are given in Appendix 5.4. The zero locus of these polynomials is $\bigcup_{i=1}^5 \mathfrak{p}_i$.

The Jacobian matrix, J_q , of $\mathfrak{p}(q)$ is given in Appendix 5.5. We examine the zero locus of the 2×2 minors of J_q to determine the multiplicity of the points in $\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3$ and the zero locus of the 3×3 minors of J_q to determine the multiplicity of the points in \mathfrak{p}_4 and \mathfrak{p}_5 [10]. The only points on which these minors vanish are $e_1, e_2 \in \mathfrak{p}_1 \cap \mathfrak{p}_2 \cap \mathfrak{p}_3$.

The coordinate ring of the point scheme is $\mathbb{k}[x_1, x_2, x_3, x_4]/I$, where I is the ideal generated by the polynomials in Appendix 5.4. Consider the plane $P = \mathcal{V}(x_2 - x_3 - x_4)$. The points of intersection of \mathfrak{p} and P are

$$\left(-\frac{q^4}{(q^4-1)^2}, 1, 0, 1\right), \left(-\frac{q^4}{(q^4-1)^2}, 1, 1, 0\right), (0, 0, 1, -1), (1, 0, 0, 0).$$

By inverting $x_1 + \frac{q^4}{(q^4-1)^2}x_3 + \frac{q^4}{(q^4-1)^2}x_4$ in the coordinate ring of $\mathfrak{p}(q) \cap P$, we can determine the multiplicity of e_1 . This yields a ring isomorphic to a polynomial ring on generators x, y with relations $x^2 = 0, xy = 0, y^2 = 0$, which is three-dimensional. Thus, e_1 is a multiple point of $\mathfrak{p}(q)$ only as a consequence of it being an intersection point of three irreducible components of $\mathfrak{p}(q)$; using the automorphism φ , we can conclude the same applies to e_2 .

Therefore, the point scheme is as proposed. \blacksquare

Corollary 3.4. *Let $A = \mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$.*

- (i) *The points in $\mathbb{P}(A_1^*) \times \mathbb{P}(A_1^*)$ on which the defining relations of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ vanish are of the form (p, p) , if $p \in \mathfrak{p}_1 \cup \mathfrak{p}_4 \cup \mathfrak{p}_5$, and, if $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \in \mathfrak{p}_j$, for $j = 2, 3$, are of the form $\left((\alpha_1, \alpha_2, \alpha_3, \alpha_4), (\alpha_1, q^{4(-1)^j}\alpha_2, q^{-2}\alpha_3, q^2\alpha_4)\right)$.*
- (ii) *There exists an automorphism $\sigma : \mathfrak{p} \rightarrow \mathfrak{p}$ which, on closed points, is defined by*

$$\sigma(p) = \sigma(p_1, p_2, p_3, p_4) = \begin{cases} (p_1, p_2, p_3, p_4), & p \in \mathfrak{p}_1 \cup \mathfrak{p}_4 \cup \mathfrak{p}_5 \\ (p_1, q^{4(-1)^j}p_2, q^{-2}p_3, q^2p_4), & p \in \mathfrak{p}_j, \text{ for } j = 2, 3. \end{cases}$$

Proof. Part (i) is shown by computation. The existence of the map in (ii) follows from (i) and [13]. \blacksquare

The quantum Casimir element of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$ is $\Omega_q = EF + FE + \left(\frac{q^4+1}{q^4-1}\right) \left(\frac{K^2+K^{-2}}{q^2-q^{-2}}\right)$ and its image in $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is $\Omega'_q = x_1x_2 + x_2x_1 + \left(\frac{q^4+1}{q^4-1}\right) \left(\frac{x_3^2+x_4^2}{q^2-q^{-2}}\right)$.

If $p \in \mathfrak{p}_2 \cup \mathfrak{p}_3$, then $\Omega'_q(p, \sigma(p)) = 0$. Moreover, Ω'_q is a central element of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$; therefore, the geometric data has identified Ω_q as a distinguished (central) element of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{k}))$.

Theorem 3.5. *The reduced variety of the line scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ is given by $\mathfrak{L}(q) = \bigcup_{i=1}^3 \mathfrak{L}_i$ where*

(I) $\mathfrak{L}_1 = \mathcal{V}(M_{13}, M_{23}, M_{34})$,

(II) $\mathfrak{L}_2 = \mathcal{V}(M_{14}, M_{24}, M_{34})$, and

(III) \mathfrak{L}_3 is given by the zero locus of the seven polynomials:

$$\begin{aligned} q^4M_{34}^2 + (q^4 - 1)^2M_{14}M_{24}, & \quad M_{12}M_{34} + M_{14}M_{23} - M_{13}M_{24}, \\ q^4M_{34}^2 + (q^4 - 1)^2M_{13}M_{23}, & \quad -M_{12}M_{24}M_{34} + M_{13}M_{24}^2 + q^4M_{23}M_{34}^2, \\ q^4(M_{24}^2 - M_{23}^2)M_{34} + (q^4 - 1)^2M_{12}M_{23}M_{24}, & \quad -M_{12}M_{13}M_{34} + M_{13}^2M_{24} + q^4M_{14}M_{34}^2, \\ q^4(M_{13}^2 - M_{14}^2)M_{34} + (q^4 - 1)^2M_{12}M_{13}M_{14}. & \end{aligned}$$

Proof. For details on the construction of the line scheme from the defining relations of a quadratic quantum \mathbb{P}^3 , the reader is referred to [4, 17].

The polynomials that define the line scheme are given in Appendix 5.6. The reduced variety given by those polynomials is $\mathfrak{L}(q)$. For details of the computation, the reader is referred to [3]. ■

The next result describes the lines in \mathbb{P}^3 that are parametrized by the closed points of the line scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$. We thank S. P. Smith of the University of Washington for his suggestion to consider a pencil of quadrics in \mathbb{P}^3 .

Corollary 3.6. *Let $\mathfrak{L}(q) = \bigcup_{i=1}^3 \mathfrak{L}_i$ be as above. The lines in \mathbb{P}^3 that correspond to the line modules of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ are precisely the lines in the pencil of quadrics*

$$Q_q(\alpha, \beta) = \mathcal{V} \left(\alpha q^4(x_3^2 + x_4^2) + \alpha(q^4 - 1)^2 x_1 x_2 + \beta q^4 x_3 x_4 \right),$$

where $(\alpha, \beta) \in \mathbb{P}^1$. More precisely,

- (i) \mathfrak{L}_1 gives all lines in $\mathcal{V}(x_3)$,
- (ii) \mathfrak{L}_2 gives all lines in $\mathcal{V}(x_4)$, and
- (iii) \mathfrak{L}_3 gives the union of the following three families of lines:
 - (a) the lines in $\mathcal{V}(x_2)$ that pass through e_1 ,
 - (b) the lines in $\mathcal{V}(x_1)$ that pass through e_2 , and
 - (c) the lines in \mathbb{P}^3 of the form $\mathcal{V}(x_1 - a_1 x_3 - b_1 x_4, x_2 - a_2 x_3 - b_2 x_4)$, such that $a_1, a_2, b_1, b_2 \in \mathbb{k}$, $q^4 + (q^4 - 1)^2 a_1 a_2 = 0$ and $a_1 a_2 = b_1 b_2$.

Proof. Let $(a_1, a_2, a_3, a_4), (b_1, b_2, b_3, b_4) \in \mathbb{P}^3$ be distinct points and let

$$\ell = \begin{bmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \end{bmatrix}$$

represent the line through them.

- (i) If ℓ is given by \mathfrak{L}_1 , then it follows that ℓ is represented by

$$\begin{bmatrix} a_1 & a_2 & 0 & a_4 \\ b_1 & b_2 & 0 & b_4 \end{bmatrix}.$$

So, ℓ belongs to $\mathcal{V}(x_3)$ if and only if ℓ is given by \mathfrak{L}_1 .

- (ii) Applying the automorphism φ to the lines described by \mathfrak{L}_1 gives that \mathfrak{L}_2 describes all lines in $\mathcal{V}(x_4)$.
- (iii) Assume that ℓ is given by \mathfrak{L}_3 .

If $M_{34} = 0$, then we may assume that

$$\ell = \begin{bmatrix} c_1 & c_2 & c_3 & c_4 \\ c_5 & c_6 & 0 & 0 \end{bmatrix},$$

where $c_1, \dots, c_6 \in \mathbb{k}$ satisfy the polynomials that define \mathfrak{L}_3 . In particular, $c_5 c_6 = 0$. If $c_6 \neq 0$, then ℓ may be represented by

$$\begin{bmatrix} c_1 & 0 & c_3 & c_4 \\ 0 & 1 & 0 & 0 \end{bmatrix},$$

where $c_1 c_3 c_4 = 0$. The cases where $c_3 = 0$ or $c_4 = 0$ are described by \mathfrak{L}_1 and \mathfrak{L}_2 . If $c_1 = 0$, then

$$\ell = \begin{bmatrix} 0 & 0 & c_3 & c_4 \\ 0 & 1 & 0 & 0 \end{bmatrix},$$

which belongs to $\mathcal{V}(x_1)$ and passes through e_2 . By a similar argument, if $c_5 \neq 0$, then ℓ belongs to $\mathcal{V}(x_2)$ and passes through e_1 .

If $M_{34} \neq 0$, we may take $M_{34} = 1$ and we may assume

$$\ell = \begin{bmatrix} a_1 & a_2 & 1 & 0 \\ b_1 & b_2 & 0 & 1 \end{bmatrix},$$

where a_1, a_2, b_1, b_2 satisfy the polynomials that define \mathfrak{L}_3 . In particular, $q^4 + (q^4 - 1)^2 a_1 a_2 = 0 = q^4 + (q^4 - 1)^2 b_1 b_2$; hence, $a_1 a_2 = b_1 b_2$. Any point on a line ℓ with this representation belongs to $\mathcal{V}(x_1 - a_1 x_3 - b_1 x_4, x_2 - a_2 x_3 - b_2 x_4)$.

The lines given by \mathfrak{L}_1 and \mathfrak{L}_2 are precisely those on $Q_q(0, 1)$. The lines given by \mathfrak{L}_3 of the form

$$\begin{bmatrix} 0 & 0 & c_3 & c_4 \\ 0 & 1 & 0 & 0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} 0 & 0 & c_3 & c_4 \\ 1 & 0 & 0 & 0 \end{bmatrix},$$

for $c_3 c_4 \neq 0$, belong to $Q_q\left(1, -\frac{c_3^2 + c_4^2}{c_3 c_4}\right)$. If $\ell = \mathcal{V}(x_1 - a_1 x_3 - b_1 x_4, x_2 - a_2 x_3 - b_2 x_4)$, where $a_1, a_2, b_1, b_2 \in \mathbb{k}$, $q^4 + (q^4 - 1)^2 a_1 a_2 = 0$ and $a_1 a_2 = b_1 b_2$, then ℓ belongs to $Q_q\left(1, -\frac{(q^4 - 1)^2}{q^4} (a_1 b_2 + a_2 b_1)\right)$.

If $\alpha \neq 0$ and $\beta^2 \neq 4$, then $Q_q(1, \beta)$ has rulings

$$\left\{ \mathcal{V} \left(x_1 - \mu q^4 (x_3 - \delta_1 x_4), \mu (q^4 - 1)^2 x_2 + x_3 - \delta_2 x_4 \right) : \mu \in \mathbb{P}^1 \right\}$$

and

$$\left\{ \mathcal{V} \left(x_2 - \mu q^4 (x_3 - \delta_1 x_4), \mu (q^4 - 1)^2 x_1 + x_3 - \delta_2 x_4 \right) : \mu \in \mathbb{P}^1 \right\},$$

where δ_1 and δ_2 are distinct solutions of $\delta^2 + \beta \delta + 1 = 0$.

If $\alpha \neq 0$ and $\beta^2 = 4$, then

$$Q_q(1, \beta) = \mathcal{V} \left(q^4 \left(x_3 + \frac{\beta}{2} x_4 \right)^2 + (q^4 - 1)^2 x_1 x_2 \right)$$

which is a rank-three quadric and so has only one ruling, namely

$$\left\{ \mathcal{V} \left(x_1 - \mu q^4 \left(x_3 + \frac{\beta}{2} x_4 \right), \mu (q^4 - 1)^2 x_2 + x_3 + \frac{\beta}{2} x_4 \right) : \mu \in \mathbb{P}^1 \right\}.$$

The lines in each of these rulings are given by \mathfrak{L}_3 . Therefore, the lines in \mathbb{P}^3 that correspond to the line modules of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ are as proposed. \blacksquare

4. THE COLOR LIE ALGEBRA $\mathfrak{sl}_k(2, \mathbb{k})$ 4.1. The \mathbb{k} -Algebra $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.

Let $\{e, f, h\}$ be the standard basis of $\mathfrak{sl}(2, \mathbb{k})$ and define

$$a_1 = \frac{i}{2}(e - f), \quad a_2 = -\frac{1}{2}(e + f), \quad a_3 = \frac{i}{2}h,$$

so that the bracket on $\mathfrak{sl}(2, \mathbb{k})$ is defined by

$$[a_1, a_2] = -a_3, \quad [a_2, a_3] = a_1, \quad [a_1, a_3] = -a_2.$$

Let $G = \mathbb{Z}_2 \times \mathbb{Z}_2$ and define a G -grading on $\mathfrak{sl}(2, \mathbb{k})$ by $\mathfrak{sl}(2, \mathbb{k}) = \bigoplus_{g \in G} X_g$, where

$$X_0 = \{0\}, \quad X_{(1,0)} = \mathbb{k}a_1, \quad X_{(0,1)} = \mathbb{k}a_2, \quad X_{(1,1)} = \mathbb{k}a_3.$$

Define a bicharacter map $\epsilon : G \times G \rightarrow \mathbb{k}^\times$ by $\epsilon((\alpha_1, \alpha_2), (\beta_1, \beta_2)) = (-1)^{\alpha_1\beta_2 - \alpha_2\beta_1}$.

The color Lie algebra $\mathfrak{sl}_k(2, \mathbb{k})$, called Klein $\mathfrak{sl}(2, \mathbb{k})$, is the ϵ -Lie algebra with bracket

$$[a_1, a_2] = a_3, \quad [a_2, a_3] = a_1, \quad [a_3, a_1] = a_2.$$

For more details on the construction of $\mathfrak{sl}_k(2, \mathbb{k})$ from $\mathfrak{sl}(2, \mathbb{k})$, the reader is referred to [5].

The universal enveloping algebra of $\mathfrak{sl}_k(2, \mathbb{k})$ is

$$\mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k})) = \frac{\mathbb{k}\langle a_1, a_2, a_3 \rangle}{\langle a_1a_2 + a_2a_1 - a_3, a_2a_3 + a_3a_2 - a_1, a_3a_1 + a_1a_3 - a_2 \rangle}.$$

Following the example of [12], we homogenize $\mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k}))$ using a central variable in order to obtain an associated quadratic quantum \mathbb{P}^3 .

Definition 4.1. The homogenization, $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$, of $\mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k}))$ is the \mathbb{k} -algebra on degree-one generators a_1, a_2, a_3, a_4 with defining relations

$$\begin{aligned} a_1a_2 + a_2a_1 &= a_3a_4, & a_2a_3 + a_3a_2 &= a_1a_4, & a_3a_1 + a_1a_3 &= a_2a_4, \\ a_1a_4 &= a_4a_1, & a_2a_4 &= a_4a_2, & a_3a_4 &= a_4a_3. \end{aligned}$$

Note that $\mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k})) \cong \mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k})) / \langle a_4 - 1 \rangle$. Unlike $\mathcal{H}(\mathfrak{sl}(1|1))$ and $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$, it is not clear if $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ is an Ore extension of an AS-regular algebra of global dimension three. We show regularity, instead, in Theorem 4.2, via a result of [13].

Theorem 4.2. *The \mathbb{k} -algebra $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ is a quadratic quantum \mathbb{P}^3 .*

Proof. A computation using Bergman's Diamond Lemma shows that a \mathbb{k} -basis for $\mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k}))$ is $\mathfrak{B} = \{a_1^{i_1} a_2^{i_2} a_3^{i_3} : i_1, i_2, i_3 = 0, 1, 2, \dots\}$. Let $D = \mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$. Note that $A = D / \langle a_4 \rangle$ is a skew-polynomial ring on three variables and therefore is an AS-regular algebra. We will prove that D is a central regular extension of A in the sense of [13].

Suppose $a_4 f = 0$ in D for some $f \in D \setminus \mathbb{k}$. We may assume $f = g_1 + g_2 + \dots + g_m \in D$ is homogeneous, where each of the g_i are scalar multiples of monomials with the generators of D

in increasing order. Let \bar{f} denote the image of f in $D/\langle a_4 - 1 \rangle$. Since $a_4 f = 0$ in D , we have $\bar{f} = 0$ in $D/\langle a_4 - 1 \rangle \cong \mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k}))$. Since the g_i are written with the generators in increasing order, it follows that an element of the form $a_1^{j_1} a_2^{j_2} a_3^{j_3}$, for some $j_1, j_2, j_3 = 0, 1, 2, \dots$, is missing from \mathfrak{B} , which is a contradiction. Hence, a_4 is regular in D .

It follows that D is a central regular extension of A and, by [13], is therefore an AS-regular algebra of global dimension four (cf. [13, Theorem 2.6, Corollary 2.7] and the paragraph after [13, Definition 3.1.1]). Moreover, A is a skew-polynomial ring, so it is Auslander-regular. Hence, by [14, Section 5.10], $D = \mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ is also Auslander-regular and satisfies the Cohen-Macaulay property. \blacksquare

We have the following automorphism on $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$. If S_4 denotes the symmetric group on $\{1, 2, 3, 4\}$ and we define $\rho = (1\ 2\ 3) \in S_4$, then the map $\varphi : \mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k})) \rightarrow \mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ defined by $\varphi(a_j) = a_{\rho(j)}$, for all j , is an automorphism of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.

4.2. The Quantum Space of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.

Recall the notation e_1, e_2, e_3, e_4 from Section 1.

Theorem 4.3. *The point scheme, \mathfrak{p} , of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ is the union of three lines and five points:*

- | | |
|--|---|
| (i) $\mathfrak{p}_1 = \mathcal{V}(a_1, a_4)$, | (v) $\mathfrak{p}_5 = \mathcal{V}(a_2 + a_1, a_3 + a_1, a_4 - 2a_1)$, |
| (ii) $\mathfrak{p}_2 = \mathcal{V}(a_2, a_4)$, | (vi) $\mathfrak{p}_6 = \mathcal{V}(a_2 + a_1, a_3 - a_1, a_4 + 2a_1)$, |
| (iii) $\mathfrak{p}_3 = \mathcal{V}(a_3, a_4)$, | (vii) $\mathfrak{p}_7 = \mathcal{V}(a_2 - a_1, a_3 - a_1, a_4 - 2a_1)$, |
| (iv) $\mathfrak{p}_4 = \mathcal{V}(a_1, a_2, a_3)$, | (viii) $\mathfrak{p}_8 = \mathcal{V}(a_2 - a_1, a_3 + a_1, a_4 + 2a_1)$, |

where the points $e_1 \in \mathfrak{p}_2 \cap \mathfrak{p}_3$, $e_2 \in \mathfrak{p}_1 \cap \mathfrak{p}_3$, $e_3 \in \mathfrak{p}_1 \cap \mathfrak{p}_2$ are counted with multiplicity three and all other points are reduced.

Proof. The polynomials that define \mathfrak{p} are listed in Appendix 5.7. The zero locus of these polynomials is $\bigcup_{i=1}^8 \mathfrak{p}_i$.

The Jacobian matrix, J , of the point scheme is given in Appendix 5.8. We examine the zero locus of the 2×2 minors of J to determine the multiplicity of the points in $\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3$, and the zero locus of the 3×3 minors of J to determine the multiplicity of the points in $\mathfrak{p}_4, \dots, \mathfrak{p}_8$ [10]. The only points where these minors vanish are the points e_1, e_2 , and e_3 .

The coordinate ring of the point scheme is $\mathbb{k}[a_1, a_2, a_3, a_4]/I$, where I is the ideal generated by the polynomials in Appendix 5.7. Consider the plane $P = \mathcal{V}(a_2 - a_3 - a_4)$. The points of intersection of \mathfrak{p} and P are e_1 and $(0, 1, 1, 0)$. We localize the coordinate ring of $\mathfrak{p} \cap P$ around e_1 and obtain an algebra that is isomorphic to a polynomial ring in one variable, x , with exactly one relation, $x^3 = 0$, and so has dimension three. Therefore, e_1 has multiplicity three; by using the automorphisms φ and φ^2 , it follows that e_2 and e_3 also have multiplicity three. \blacksquare

Corollary 4.4. *Let $A = \mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.*

- (i) *The points in $\mathbb{P}(A_1^*) \times \mathbb{P}(A_1^*)$ on which the defining relations of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ vanish are of the form (p, p) , if $p \in \bigcup_{i=4}^8 \mathfrak{p}_i$, and are of the form*

$$\left((\alpha_1, \alpha_2, \alpha_3, 0), ((-1)^{\delta_{j2}}\alpha_1, (-1)^{\delta_{j3}}\alpha_2, (-1)^{\delta_{j1}}\alpha_3, 0) \right),$$

if $(\alpha_1, \alpha_2, \alpha_3, 0) \in \mathfrak{p}_j$, for $j = 1, 2, 3$, where δ_{jk} is the Kronecker-delta symbol.

- (ii) *There exists an automorphism $\sigma : \mathfrak{p} \rightarrow \mathfrak{p}$ which, on closed points, is defined by*

$$\sigma(p) = \sigma(p_1, p_2, p_3, p_4) = \begin{cases} (p_1, p_2, p_3, p_4), & p \in \mathfrak{p}_j, j = 4, \dots, 8 \\ ((-1)^{\delta_{j2}}p_1, (-1)^{\delta_{j3}}p_2, (-1)^{\delta_{j1}}p_3, 0), & p \in \mathfrak{p}_j, j = 1, 2, 3. \end{cases}$$

Proof. Part (i) is shown by computation. The existence of the map in (ii) follows from (i) and [13]. ■

Theorem 4.5. *The line scheme of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ has dimension two and its reduced variety \mathfrak{L} is the union of twelve lines and one plane:*

- (I) $\mathfrak{L}_0 = \mathcal{V}(M_{14}, M_{24}, M_{34})$,
- (II) $\mathfrak{L}_1 = \mathcal{V}(M_{12}, M_{34}, M_{14} - M_{24}, M_{13} - M_{23})$,
- (III) $\mathfrak{L}_2 = \mathcal{V}(M_{12}, M_{34}, M_{14} + M_{24}, M_{13} + M_{23})$,
- (IV) $\mathfrak{L}_3 = \mathcal{V}(M_{34}, M_{14} + 2M_{23}, 2M_{13} + M_{24}, 2M_{23} - M_{24})$,
- (V) $\mathfrak{L}_4 = \mathcal{V}(M_{34}, M_{14} + 2M_{23}, 2M_{13} + M_{24}, 2M_{23} + M_{24})$,
- (VI) $\mathfrak{L}_5 = \mathcal{V}(M_{13}, M_{24}, M_{34} - M_{14}, M_{12} + M_{23})$,
- (VII) $\mathfrak{L}_6 = \mathcal{V}(M_{13}, M_{24}, M_{34} + M_{14}, M_{12} - M_{23})$,
- (VIII) $\mathfrak{L}_7 = \mathcal{V}(M_{24}, M_{34} + 2M_{12}, M_{14} - 2M_{23}, 2M_{12} - M_{14})$,
- (IX) $\mathfrak{L}_8 = \mathcal{V}(M_{24}, M_{34} + 2M_{12}, M_{14} - 2M_{23}, 2M_{12} + M_{14})$,
- (X) $\mathfrak{L}_9 = \mathcal{V}(M_{14}, M_{23}, M_{24} - M_{34}, M_{13} - M_{12})$,
- (XI) $\mathfrak{L}_{10} = \mathcal{V}(M_{14}, M_{23}, M_{24} + M_{34}, M_{13} + M_{12})$,
- (XII) $\mathfrak{L}_{11} = \mathcal{V}(M_{14}, M_{24} - 2M_{13}, M_{34} - 2M_{12}, M_{34} + 2M_{13})$,
- (XIII) $\mathfrak{L}_{12} = \mathcal{V}(M_{14}, M_{24} - 2M_{13}, M_{34} - 2M_{12}, M_{34} - 2M_{13})$.

Proof. For details on the construction of the line scheme from the defining relations of a quadratic quantum \mathbb{P}^3 , the reader is referred to [4, 17].

The polynomials that define the line scheme are given in Appendix 5.9. The reduced variety given by those polynomials is \mathfrak{L} . For details of the computation, the reader is referred to [3]. ■

Corollary 4.6. *The lines in \mathbb{P}^3 that are parametrized by the line scheme of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ are:*

- (i) *all lines in $\mathcal{V}(a_4)$,*
- (ii) *those lines in $\mathcal{V}(a_1 - a_2)$ that pass through $(1, 1, 0, 0)$,*
- (iii) *those lines in $\mathcal{V}(a_1 + a_2)$ that pass through $(1, -1, 0, 0)$,*

- (iv) *those lines in $\mathcal{V}(a_4 - 2a_3)$ that pass through $(1, -1, 0, 0)$,*
- (v) *those lines in $\mathcal{V}(a_4 + 2a_3)$ that pass through $(1, 1, 0, 0)$,*
- (vi) *those lines in $\mathcal{V}(a_1 - a_3)$ that pass through $(1, 0, 1, 0)$,*
- (vii) *those lines in $\mathcal{V}(a_1 + a_3)$ that pass through $(1, 0, -1, 0)$,*
- (viii) *those lines in $\mathcal{V}(a_4 - 2a_2)$ that pass through $(1, 0, -1, 0)$,*
- (ix) *those lines in $\mathcal{V}(a_4 + 2a_2)$ that pass through $(1, 0, 1, 0)$,*
- (x) *those lines in $\mathcal{V}(a_2 - a_3)$ that pass through $(0, 1, 1, 0)$,*
- (xi) *those lines in $\mathcal{V}(a_2 + a_3)$ that pass through $(0, 1, -1, 0)$,*
- (xii) *those lines in $\mathcal{V}(a_4 - 2a_1)$ that pass through $(0, 1, -1, 0)$, and*
- (xiii) *those lines in $\mathcal{V}(a_4 + 2a_1)$ that pass through $(0, 1, 1, 0)$.*

Proof. Let $(b_1, b_2, b_3, b_4), (c_1, c_2, c_3, c_4) \in \mathbb{P}^3$ be distinct points and let

$$\ell = \begin{bmatrix} b_1 & b_2 & b_3 & b_4 \\ c_1 & c_2 & c_3 & c_4 \end{bmatrix}$$

represent the line through them.

- (i) If ℓ is given by \mathfrak{L}_0 , then it follows that ℓ is given by

$$\begin{bmatrix} b_1 & b_2 & b_3 & 0 \\ c_1 & c_2 & c_3 & 0 \end{bmatrix}.$$

So, ℓ is given by \mathfrak{L}_0 if and only if ℓ belongs to $\mathcal{V}(a_4)$.

- (ii) If ℓ is given by \mathfrak{L}_1 , then it follows that ℓ may be represented as

$$\begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & d_1 & d_2 \end{bmatrix},$$

where $(d_1, d_2) \in \mathbb{P}^1$. Therefore, ℓ passes through $(1, 1, 0, 0)$ and belongs to the plane $\mathcal{V}(a_1 - a_2)$.

- (iii) The result follows by an argument similar to that in (ii).

- (iv) If ℓ is given by \mathfrak{L}_3 , then

$$\ell = \begin{bmatrix} d_1 & d_2 & d_3 & d_4 \\ d_5 & d_6 & 0 & 0 \end{bmatrix},$$

where $d_1, \dots, d_6 \in \mathbb{k}$ satisfy the polynomials that define \mathfrak{L}_3 .

If $d_6 = 0$, then it follows that ℓ may be represented as

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix},$$

which is the line $\mathcal{V}(a_3, a_4)$ (this line is also given by \mathfrak{L}_0).

If $d_6 \neq 0$, then it follows that $\ell = \mathcal{V}(a_3, a_4)$ or

$$\ell = \begin{bmatrix} 0 & d_7 & d_3 & 2d_3 \\ 1 & -1 & 0 & 0 \end{bmatrix},$$

where $d_7 \in \mathbb{k}$. Therefore, \mathfrak{L}_3 gives all lines that pass through $(1, -1, 0, 0)$ and belong to $\mathcal{V}(a_4 - 2a_3)$.

(v) The result follows by an argument similar to that in (iv).

To obtain the lines given by $\mathfrak{L}_5, \dots, \mathfrak{L}_8$, one may apply φ^2 to the lines described by $\mathfrak{L}_1, \dots, \mathfrak{L}_4$, respectively. To obtain the lines given by $\mathfrak{L}_9, \dots, \mathfrak{L}_{12}$, one may apply φ to the lines described by $\mathfrak{L}_1, \dots, \mathfrak{L}_4$, respectively. \blacksquare

We conclude by remarking that, unlike $\mathcal{H}(\mathfrak{sl}(2, \mathbb{k}))$, $\mathcal{H}(\mathfrak{sl}(1|1))$ or $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$, the quantum space of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ does not contain a distinguished conic or quadric. In regards to the other algebras, certain conics and quadrics in the point schemes identified distinguished elements of the algebra, including an analogue of a Casimir element for each associated enveloping algebra. This suggests that $\mathcal{U}(\mathfrak{sl}_k(2, \mathbb{k}))$ lacks such an element.

5. APPENDIX

5.1. The Polynomials Defining the Point Scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$.

$$5.1.1. \quad x_4^2(2x_1x_2 - x_3x_4)$$

$$5.1.2. \quad x_3x_4(2x_1x_2 - x_3x_4)$$

$$5.1.3. \quad x_3^2(2x_1x_2 - x_3x_4)$$

$$5.1.4. \quad x_2x_4(2x_1x_2 - x_3x_4)$$

$$5.1.5. \quad x_2x_3(2x_1x_2 - x_3x_4)$$

$$5.1.6. \quad x_1x_4(2x_1x_2 - x_3x_4)$$

$$5.1.7. \quad x_1x_3(2x_1x_2 - x_3x_4)$$

5.2. The Jacobian Matrix of the Point Scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$.

$$\begin{bmatrix} 2x_2x_4^2 & 2x_1x_4^2 & -x_4^3 & 4x_1x_2x_4 - 3x_3x_4^2 \\ 2x_2x_3x_4 & 2x_1x_3x_4 & 2x_1x_2x_4 - 2x_3x_4^2 & 2x_1x_2x_3 - 2x_3^2x_4 \\ 2x_2x_3^2 & 2x_1x_3^2 & 4x_1x_2x_3 - 3x_3^2x_4 & -x_3^3 \\ 2x_2^2x_4 & 4x_1x_2x_4 - x_3x_4^2 & -x_2x_4^2 & 2x_1x_2^2 - 2x_2x_3x_4 \\ 2x_2^2x_3 & 4x_1x_2x_3 - x_3^2x_4 & 2x_1x_2^2 - 2x_2x_3x_4 & -x_2x_3^2 \\ 4x_1x_2x_4 - x_3x_4^2 & 2x_1^2x_4 & -x_1x_4^2 & 2x_1^2x_2 - 2x_1x_3x_4 \\ 4x_1x_2x_3 - x_3^2x_4 & 2x_1^2x_3 & 2x_1^2x_2 - 2x_1x_3x_4 & -x_1x_3^2 \end{bmatrix}.$$

5.3. The Polynomials Defining the Line Scheme of $\mathcal{H}(\mathfrak{sl}(1|1))$.

- 5.3.1. $-M_{13}M_{14}(M_{13}M_{24} - M_{14}M_{23})$
- 5.3.2. $(M_{14}M_{23} - M_{13}M_{24})(2M_{12}^2 + M_{13}M_{24} + M_{14}M_{23})$
- 5.3.3. $M_{23}M_{24}(M_{14}M_{23} - M_{13}M_{24})$
- 5.3.4. $M_{12}M_{13}M_{14}M_{34}$
- 5.3.5. $M_{13}^2M_{14}M_{34}$
- 5.3.6. $-M_{13}M_{14}^2M_{34}$
- 5.3.7. $-2M_{12}M_{13}M_{23}M_{34}$
- 5.3.8. $-2M_{13}^2M_{23}M_{34}$
- 5.3.9. $-M_{13}M_{14}M_{23}M_{34}$
- 5.3.10. $-2M_{13}M_{23}^2M_{34}$
- 5.3.11. $-2M_{12}M_{14}M_{24}M_{34}$
- 5.3.12. $M_{13}M_{14}M_{24}M_{34}$
- 5.3.13. $-2M_{14}^2M_{24}M_{34}$
- 5.3.14. $M_{12}M_{23}M_{24}M_{34}$
- 5.3.15. $-M_{13}M_{23}M_{24}M_{34}$
- 5.3.16. $M_{13}M_{23}M_{24}M_{34}$
- 5.3.17. $-M_{14}M_{23}M_{24}M_{34}$
- 5.3.18. $M_{14}M_{23}M_{24}M_{34}$
- 5.3.19. $-M_{23}^2M_{24}M_{34}$
- 5.3.20. $-2M_{14}M_{24}^2M_{34}$
- 5.3.21. $M_{23}M_{24}^2M_{34}$
- 5.3.22. $-M_{13}M_{14}M_{34}^2$
- 5.3.23. $-M_{23}M_{24}M_{34}^2$
- 5.3.24. $M_{12}M_{34} - M_{13}M_{24} + M_{14}M_{23}$
- 5.3.25. $-M_{14}(M_{12}^2M_{34} + M_{12}M_{13}M_{24} - M_{12}M_{14}M_{23} + M_{14}M_{23}M_{34})$
- 5.3.26. $M_{23}(M_{12}^2M_{34} + M_{12}M_{13}M_{24} - M_{12}M_{14}M_{23} + M_{14}M_{23}M_{34})$
- 5.3.27. $M_{13}(M_{12}^2M_{34} + M_{12}M_{13}M_{24} - M_{12}M_{14}M_{23} + M_{13}M_{24}M_{34})$
- 5.3.28. $-M_{24}(M_{12}^2M_{34} + M_{12}M_{13}M_{24} - M_{12}M_{14}M_{23} + M_{13}M_{24}M_{34})$
- 5.3.29. $M_{13}^2M_{34}(2M_{12} + M_{34})$
- 5.3.30. $-M_{14}^2M_{34}(2M_{12} - M_{34})$

- 5.3.31. $-M_{13}M_{23}(-M_{12}M_{34} + M_{13}M_{24} - M_{14}M_{23} + M_{34}^2)$
 5.3.32. $-M_{13}M_{23}(-M_{12}M_{34} + M_{13}M_{24} - M_{14}M_{23} - M_{34}^2)$
 5.3.33. $M_{14}M_{23}M_{34}(2M_{12} - M_{34})$
 5.3.34. $M_{23}^2M_{34}(2M_{12} - M_{34})$
 5.3.35. $M_{13}M_{24}M_{34}(2M_{12} + M_{34})$
 5.3.36. $-M_{14}M_{24}(M_{12}M_{34} - M_{13}M_{24} + M_{14}M_{23} + M_{34}^2)$
 5.3.37. $-M_{14}M_{24}(M_{12}M_{34} - M_{13}M_{24} + M_{14}M_{23} - M_{34}^2)$
 5.3.38. $-M_{24}^2M_{34}(2M_{12} + M_{34})$
 5.3.39. $M_{13}M_{34}(M_{12}M_{34} + M_{13}M_{24} + M_{14}M_{23} + M_{34}^2)$
 5.3.40. $M_{14}M_{34}(-M_{12}M_{34} + M_{13}M_{24} + M_{14}M_{23} + M_{34}^2)$
 5.3.41. $M_{23}M_{34}(-M_{12}M_{34} + M_{13}M_{24} + M_{14}M_{23} + M_{34}^2)$
 5.3.42. $M_{24}M_{34}(M_{12}M_{34} + M_{13}M_{24} + M_{14}M_{23} + M_{34}^2)$
 5.3.43. $M_{34}^2(2M_{13}M_{24} + 2M_{14}M_{23} + M_{34}^2)$

5.4. The Polynomials Defining the Point Scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$.

- 5.4.1. $-(q^4 - 1)x_1x_3^2x_4$
 5.4.2. $q^{-4}(q^4 - 1)x_2x_3^2x_4$
 5.4.3. $-q^{-2}(q^4 - 1)^{-1}x_3^2(q^8x_1x_2 - 2q^4x_1x_2 + q^4x_3^2 - q^4x_4^2 + x_1x_2)$
 5.4.4. $-q^{-2}(q^4 - 1)x_1x_3x_4^2$
 5.4.5. $(q^4 - 1)x_1^2x_3x_4$
 5.4.6. $(q^4 - 1)^{-1}x_3x_4(x_3 - x_4)(x_3 + x_4)$
 5.4.7. $q^{-2}(q^4 - 1)x_2x_3x_4^2$
 5.4.8. $-(q^4 - 1)^{-1}q^4x_3x_4(x_3 - x_4)(x_3 + x_4)$
 5.4.9. $-q^{-4}(q^4 - 1)x_2^2x_3x_4$
 5.4.10. $q^{-2}(q^4 - 1)^{-1}x_4^2((q^4 - 1)^2x_1x_2 - q^4x_3^2 + q^4x_4^2)$
 5.4.11. $-q^{-4}(q^4 - 1)(q^4 + 1)x_1x_2x_3x_4$
 5.4.12. $q^{-2}(q^4 - 1)^{-1}x_1x_3((q^4 - 1)^2x_1x_2 - q^8x_4^2 + q^4x_3^2)$
 5.4.13. $q^{-2}(q^4 - 1)^{-1}x_2x_3((q^4 - 1)^2x_1x_2 + q^4x_3^2 - x_4^2)$
 5.4.14. $-(q^4 - 1)^{-1}x_1x_4((q^4 - 1)^2x_1x_2 + q^4x_4^2 - x_3^2)$
 5.4.15. $-q^{-4}(q^4 - 1)^{-1}x_2x_4((q^4 - 1)^2x_1x_2 - q^8x_3^2 + q^4x_4^2)$

5.5. The Jacobian Matrix of the Point Scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$.

For typesetting purposes, we display the Jacobian matrix associated to the point scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$ as two submatrices: the first matrix displays the first two columns of the Jacobian matrix while the second matrix displays the third and fourth columns:

$$\left[\begin{array}{cc} (1 - q^4) x_3^2 x_4 & 0 \\ 0 & q^{-4} (q^4 - 1) x_3^2 x_4 \\ -q^{-2} (q^4 - 1) x_2 x_3^2 & -q^{-2} (q^4 - 1) x_1 x_3^2 \\ -q^{-2} (q^4 - 1) x_3 x_4^2 & 0 \\ 2 (q^4 - 1) x_1 x_3 x_4 & 0 \\ 0 & 0 \\ 0 & q^{-2} (q^4 - 1) x_3 x_4^2 \\ 0 & 0 \\ 0 & -2q^{-4} (q^4 - 1) x_2 x_3 x_4 \\ q^{-2} (q^4 - 1) x_2 x_4^2 & q^{-2} (q^4 - 1) x_1 x_4^2 \\ -q^{-4} (q^4 - 1) (q^4 + 1) x_2 x_3 x_4 & -q^{-4} (q^4 - 1) (q^4 + 1) x_1 x_3 x_4 \\ q^{-2} (q^4 - 1)^{-1} x_3 (2(q^4 - 1)^2 x_1 x_2 + q^4 x_3^2 - q^8 x_4^2) & q^{-2} (q^4 - 1) x_1^2 x_3 \\ q^{-2} (q^4 - 1) x_2^2 x_3 & q^{-2} (q^4 - 1)^{-1} x_3 (2(q^4 - 1)^2 x_1 x_2 + q^4 x_3^2 - x_4^2) \\ -(q^4 - 1)^{-1} x_4 (2 (q^4 - 1)^2 x_1 x_2 + q^4 x_4^2 - x_3^2) & (1 - q^4) x_1^2 x_4 \\ -q^{-4} (q^4 - 1) x_2^2 x_4 & -q^{-4} (q^4 - 1)^{-1} x_4 (2 (q^4 - 1)^2 x_1 x_2 - q^8 x_3^2 + q^4 x_4^2) \end{array} \right]$$

$$\left[\begin{array}{cc} -2 (q^4 - 1) x_1 x_3 x_4 & (1 - q^4) x_1 x_3^2 \\ 2q^{-4} (q^4 - 1) x_2 x_3 x_4 & q^{-4} (q^4 - 1) x_2 x_3^2 \\ -2q^{-2} (q^4 - 1)^{-1} x_3 ((q^4 - 1)^2 x_1 x_2 + 2q^4 x_3^2 - q^4 x_4^2) & 2q^2 (q^4 - 1)^{-1} x_3^2 x_4 \\ -q^{-2} (q^4 - 1) x_1 x_4^2 & -2q^{-2} (q^4 - 1) x_1 x_3 x_4 \\ (q^4 - 1) x_1^2 x_4 & (q^4 - 1) x_1^2 x_3 \\ (q^4 - 1)^{-1} x_4 (3x_3^2 - x_4^2) & (q^4 - 1)^{-1} x_3 (x_3^2 - 3x_4^2) \\ q^{-2} (q^4 - 1) x_2 x_4^2 & 2q^{-2} (q^4 - 1) x_2 x_3 x_4 \\ -(q^4 - 1)^{-1} q^4 x_4 (3x_3^2 - x_4^2) & -(q^4 - 1)^{-1} q^4 x_3 (x_3^2 - 3x_4^2) \\ -q^{-4} (q^4 - 1) x_2^2 x_4 & -q^{-4} (q^4 - 1) x_2^2 x_3 \\ -2q^2 (q^4 - 1)^{-1} x_3 x_4^2 & 2q^{-1} (q^4 - 1)^{-1} x_4 ((q^4 - 1)^2 x_1 x_2 - q^4 x_3^2 + 2q^4 x_4^2) \\ -q^{-4} (q - 1)(q + 1) (q^2 + 1) (q^4 + 1) x_1 x_2 x_4 & -q^{-4} (q - 1)(q + 1) (q^2 + 1) (q^4 + 1) x_1 x_2 x_3 \\ q^{-2} (q^4 - 1)^{-1} x_1 ((q^4 - 1)^2 x_1 x_2 - q^8 x_4^2 + 3q^4 x_3^2) & -2(q^4 - 1)^{-1} q^6 x_1 x_3 x_4 \\ q^{-2} (q^4 - 1)^{-1} x_2 ((q^4 - 1)^2 x_1 x_2 + 3q^4 x_3^2 - x_4^2) & -2q^{-2} (q^4 - 1)^{-1} x_2 x_3 x_4 \\ 2(q^4 - 1)^{-1} x_1 x_3 x_4 & -(q^4 - 1)^{-1} x_1 ((q^4 - 1)^2 x_1 x_2 + 3q^4 x_4^2 - x_3^2) \\ 2q^4 (q^4 - 1)^{-1} x_2 x_3 x_4 & -q^{-4} (q^4 - 1)^{-1} x_2 ((q^4 - 1)^2 x_1 x_2 - q^8 x_3^2 + 3q^4 x_4^2) \end{array} \right]$$

5.6. The Polynomials Defining the Line Scheme of $\mathcal{H}_q(\mathfrak{sl}(2, \mathbb{k}))$.

- 5.6.1. $M_{14}M_{23} - M_{13}M_{24} + M_{12}M_{34}$
- 5.6.2. $M_{13}(M_{12}M_{13}M_{14}q^8 - 2M_{12}M_{13}M_{14}q^4 + M_{13}^2M_{34}q^4 - M_{14}^2M_{34}q^4 + M_{12}M_{13}M_{14})$
- 5.6.3. $M_{14}(M_{12}M_{23}M_{24}q^8 - 2M_{12}M_{23}M_{24}q^4 - M_{23}^2M_{34}q^4 + M_{24}^2M_{34}q^4 + M_{12}M_{23}M_{24})$
- 5.6.4. $M_{24}(M_{12}M_{23}M_{24}q^8 - 2M_{12}M_{23}M_{24}q^4 - M_{23}^2M_{34}q^4 + M_{24}^2M_{34}q^4 + M_{12}M_{23}M_{24})$
- 5.6.5. $M_{12}(M_{12}M_{23}M_{24}q^8 - 2M_{12}M_{23}M_{24}q^4 - M_{23}^2M_{34}q^4 + M_{24}^2M_{34}q^4 + M_{12}M_{23}M_{24})$
- 5.6.6. $M_{34}(M_{12}M_{13}M_{14}q^8 - 2M_{12}M_{13}M_{14}q^4 + M_{13}^2M_{34}q^4 - M_{14}^2M_{34}q^4 + M_{12}M_{13}M_{14})$
- 5.6.7. $(M_{13}M_{23} - M_{14}M_{24})M_{34}^2q^2$
- 5.6.8. $(M_{13}M_{23} - M_{14}M_{24})M_{34}^2q^2$
- 5.6.9. $M_{34}(M_{12}M_{23}M_{24}q^8 - 2M_{12}M_{23}M_{24}q^4 - M_{23}^2M_{34}q^4 + M_{24}^2M_{34}q^4 + M_{12}M_{23}M_{24})$
- 5.6.10. $M_{12}M_{34}(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.11. $M_{12}M_{34}(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.12. $M_{13}M_{34}(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.13. $M_{34}(-M_{13}M_{14}M_{24}q^8 - M_{13}M_{34}^2q^4 + M_{14}^2M_{23}q^4 + M_{13}M_{14}M_{24}q^4 + M_{12}M_{14}M_{34}q^4 - M_{14}^2M_{23} - M_{12}M_{14}M_{34})$
- 5.6.14. $M_{13}M_{34}(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.15. $M_{14}M_{34}(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.16. $M_{34}(M_{13}^2M_{24}q^8 - M_{12}M_{13}M_{34}q^8 + M_{14}M_{34}^2q^4 - M_{13}M_{14}M_{23}q^4 - M_{13}^2M_{24}q^4 + M_{12}M_{13}M_{34}q^4 + M_{13}M_{14}M_{23})$
- 5.6.17. $M_{14}M_{34}(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.18. $M_{23}M_{34}(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.19. $M_{23}M_{34}(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.20. $M_{34}(-M_{13}M_{24}^2q^8 + M_{12}M_{24}M_{34}q^8 + M_{13}M_{24}^2q^4 - M_{23}M_{34}^2q^4 + M_{14}M_{23}M_{24}q^4 - M_{12}M_{24}M_{34}q^4 - M_{14}M_{23}M_{24})$
- 5.6.21. $M_{24}M_{34}(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.22. $M_{34}(-M_{13}M_{23}M_{24}q^8 + M_{14}M_{23}^2q^4 - M_{24}M_{34}^2q^4 + M_{13}M_{23}M_{24}q^4 + M_{12}M_{23}M_{34}q^4 - M_{14}M_{23}^2 - M_{12}M_{23}M_{34})$
- 5.6.23. $M_{24}M_{34}(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.24. $M_{34}^2(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.25. $M_{34}^2(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.26. $M_{14}(M_{12}M_{13}M_{14}q^8 - 2M_{12}M_{13}M_{14}q^4 + M_{13}^2M_{34}q^4 - M_{14}^2M_{34}q^4 + M_{12}M_{13}M_{14})$
- 5.6.27. $M_{12}M_{13}M_{14}M_{23}q^8 + M_{13}^2M_{23}M_{34}q^8 - M_{13}M_{14}M_{24}M_{34}q^8 + M_{12}M_{14}M_{34}^2q^4 - 2M_{12}M_{13}M_{14}M_{23}q^4 + M_{12}M_{13}M_{14}M_{23}$

- 5.6.28. $M_{12}M_{14}^2M_{23}q^8 + M_{12}^2M_{14}M_{34}q^8 - M_{13}M_{14}M_{23}M_{34}q^8 + M_{14}^2M_{24}M_{34}q^8$
 $+ M_{12}M_{13}M_{34}^2q^4 - M_{12}M_{14}^2M_{23}q^4 - M_{12}M_{13}M_{14}M_{24}q^4 - M_{12}^2M_{14}M_{34}q^4$
 $+ M_{12}M_{13}M_{14}M_{24}$
- 5.6.29. $M_{12}M_{14}M_{23}^2q^8 + M_{13}M_{23}^2M_{34}q^8 + M_{12}^2M_{23}M_{34}q^8 - M_{14}M_{23}M_{24}M_{34}q^8$
 $- M_{12}M_{14}M_{23}^2q^4 + M_{12}M_{24}M_{34}^2q^4 - M_{12}M_{13}M_{23}M_{24}q^4 - M_{12}^2M_{23}M_{34}q^4$
 $+ M_{12}M_{13}M_{23}M_{24}$
- 5.6.30. $M_{13}(M_{12}M_{23}M_{24}q^8 - 2M_{12}M_{23}M_{24}q^4 - M_{23}^2M_{34}q^4 + M_{24}^2M_{34}q^4 + M_{12}M_{23}M_{24})$
- 5.6.31. $M_{23}(M_{12}M_{23}M_{24}q^8 - 2M_{12}M_{23}M_{24}q^4 - M_{23}^2M_{34}q^4 + M_{24}^2M_{34}q^4 + M_{12}M_{23}M_{24})$
- 5.6.32. $M_{12}M_{13}M_{14}M_{24}q^8 + M_{12}M_{13}M_{34}^2q^4 - 2M_{12}M_{13}M_{14}M_{24}q^4 + M_{12}M_{13}M_{14}M_{24}$
 $+ M_{13}M_{14}M_{23}M_{34} - M_{14}^2M_{24}M_{34}$
- 5.6.33. $M_{12}M_{13}M_{14}M_{23}q^8 + M_{12}M_{14}M_{34}^2q^4 - M_{12}M_{13}M_{14}M_{23}q^4 - M_{12}M_{13}^2M_{24}q^4$
 $+ M_{12}^2M_{13}M_{34}q^4 + M_{12}M_{13}^2M_{24} - M_{12}^2M_{13}M_{34} - M_{13}^2M_{23}M_{34}$
 $+ M_{13}M_{14}M_{24}M_{34}$
- 5.6.34. $M_{13}M_{14}(q^4 + 1)(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.35. $M_{13}M_{14}(q^4 + 1)(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.36. $M_{13}M_{14}^2M_{23}q^{12} + M_{14}^2M_{34}^2q^8 - M_{13}M_{14}^2M_{23}q^8 + M_{12}M_{13}M_{14}M_{34}q^8 + M_{13}^2M_{34}^2q^4 -$
 $M_{13}^2M_{14}M_{24}q^4 - M_{12}M_{13}M_{14}M_{34}q^4 + M_{13}^2M_{14}M_{24}$
- 5.6.37. $M_{12}M_{14}M_{23}M_{24}q^8 - M_{12}M_{13}M_{24}^2q^4 + M_{12}M_{23}M_{34}^2q^4 - M_{12}M_{14}M_{23}M_{24}q^4$
 $+ M_{12}^2M_{24}M_{34}q^4 + M_{12}M_{13}M_{24}^2 - M_{14}M_{24}^2M_{34} - M_{12}^2M_{24}M_{34}$
 $+ M_{13}M_{23}M_{24}M_{34}$
- 5.6.38. $M_{13}M_{14}M_{23}M_{24}q^{12} + M_{13}M_{23}M_{34}^2q^8 - M_{13}M_{14}M_{23}M_{24}q^8 + M_{14}M_{24}M_{34}^2q^4$
 $- M_{13}M_{14}M_{23}M_{24}q^4 + M_{13}M_{14}M_{23}M_{24}$
- 5.6.39. $M_{14}M_{23}^2M_{24}q^{12} + M_{23}^2M_{34}^2q^8 - M_{14}M_{23}^2M_{24}q^8 + M_{12}M_{23}M_{24}M_{34}q^8 - M_{13}M_{23}M_{24}^2q^4 +$
 $M_{24}^2M_{34}^2q^4 - M_{12}M_{23}M_{24}M_{34}q^4 + M_{13}M_{23}M_{24}^2$
- 5.6.40. $-M_{13}^2M_{23}M_{24}q^{12} - M_{13}M_{24}M_{34}^2q^8 + M_{13}^2M_{23}M_{24}q^8 + M_{12}M_{13}M_{23}M_{34}q^8$
 $+ M_{13}M_{14}M_{23}^2q^4 - M_{14}M_{23}M_{34}^2q^4 - M_{12}M_{13}M_{23}M_{34}q^4 - M_{13}M_{14}M_{23}^2$
- 5.6.41. $-M_{13}M_{14}M_{24}^2q^{12} + M_{13}M_{14}M_{24}^2q^8 - M_{13}M_{24}M_{34}^2q^8 + M_{12}M_{14}M_{24}M_{34}q^8$
 $- M_{14}M_{23}M_{34}^2q^4 + M_{14}^2M_{23}M_{24}q^4 - M_{12}M_{14}M_{24}M_{34}q^4 - M_{14}^2M_{23}M_{24}$
- 5.6.42. $M_{23}M_{24}(q^4 + 1)(M_{13}M_{23}q^8 + M_{34}^2q^4 - 2M_{13}M_{23}q^4 + M_{13}M_{23})$
- 5.6.43. $M_{23}M_{24}(q^4 + 1)(M_{14}M_{24}q^8 + M_{34}^2q^4 - 2M_{14}M_{24}q^4 + M_{14}M_{24})$
- 5.6.44. $M_{12}^2M_{13}M_{14}q^8 - M_{13}^2M_{14}M_{23}q^8 + M_{13}M_{14}^2M_{24}q^8 - 2M_{12}^2M_{13}M_{14}q^4 + M_{12}M_{13}^2M_{34}q^4 -$
 $M_{12}M_{14}^2M_{34}q^4 + M_{12}^2M_{13}M_{14} + M_{13}^2M_{14}M_{23} - M_{13}M_{14}^2M_{24}$
- 5.6.45. $(M_{13}M_{23} - M_{14}M_{24})(-M_{14}M_{23}q^8 + M_{12}M_{34}q^4 + M_{13}M_{24})$

5.7. The Polynomials Defining the Point Scheme of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.

- 5.7.1. $a_2(a_4a_1^2 - 2a_2a_3a_1 - a_2^2a_4 + a_3^2a_4)$
- 5.7.2. $-a_1(a_4a_1^2 + 2a_2a_3a_1 - a_2^2a_4 - a_3^2a_4)$
- 5.7.3. $a_3(a_4a_1^2 - 2a_2a_3a_1 + a_2^2a_4 - a_3^2a_4)$
- 5.7.4. $a_1a_4(2a_1a_2 - a_3a_4)$
- 5.7.5. $-(a_2 - a_3)(a_2 + a_3)a_4^2$
- 5.7.6. $-a_1a_4(2a_1a_3 - a_2a_4)$
- 5.7.7. $a_2a_4(2a_1a_2 - a_3a_4)$
- 5.7.8. $a_2a_4(2a_2a_3 - a_1a_4)$
- 5.7.9. $(a_1 - a_3)(a_1 + a_3)a_4^2$
- 5.7.10. $-a_4^2(2a_1a_2 - a_3a_4)$
- 5.7.11. $(a_1 - a_2)(a_1 + a_2)a_4^2$
- 5.7.12. $a_3a_4(2a_2a_3 - a_1a_4)$
- 5.7.13. $a_3a_4(2a_1a_3 - a_2a_4)$
- 5.7.14. $-a_4^2(2a_1a_3 - a_2a_4)$
- 5.7.15. $-a_4^2(2a_2a_3 - a_1a_4)$

5.8. The Jacobian Matrix of the Point Scheme of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.

For typesetting purposes, we display the Jacobian matrix associated to the point scheme of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$ as two submatrices: the first matrix displays the first two columns of the Jacobian matrix while the second matrix displays the third and fourth columns:

$$\left[\begin{array}{cc} -2a_2(a_2a_3 - a_1a_4) & a_1^2a_4 - 4a_1a_2a_3 - 3a_2^2a_4 + a_3^2a_4 \\ -3a_1^2a_4 - 4a_1a_2a_3 + a_2^2a_4 + a_3^2a_4 & -2a_1(a_1a_3 - a_2a_4) \\ -2a_3(a_2a_3 - a_1a_4) & -2a_3(a_1a_3 - a_2a_4) \\ a_4(4a_1a_2 - a_3a_4) & 2a_1^2a_4 \\ 0 & -2a_2a_4^2 \\ -a_4(4a_1a_3 - a_2a_4) & a_1a_4^2 \\ 2a_2^2a_4 & a_4(4a_1a_2 - a_3a_4) \\ -a_2a_4^2 & a_4(4a_2a_3 - a_1a_4) \\ 2a_1a_4^2 & 0 \\ -2a_2a_4^2 & -2a_1a_4^2 \\ 2a_1a_4^2 & -2a_2a_4^2 \\ -a_3a_4^2 & 2a_3^2a_4 \\ 2a_3^2a_4 & -a_3a_4^2 \\ -2a_3a_4^2 & a_4^3 \\ a_4^3 & -2a_3a_4^2 \end{array} \right]$$

$$\begin{bmatrix} -2a_2(a_1a_2 - a_3a_4) & a_2(a_1^2 - a_2^2 + a_3^2) \\ -2a_1(a_1a_2 - a_3a_4) & -a_1(a_1^2 - a_2^2 - a_3^2) \\ a_1^2a_4 - 4a_1a_2a_3 + a_2^2a_4 - 3a_3^2a_4 & a_3(a_1^2 + a_2^2 - a_3^2) \\ -a_1a_4^2 & 2a_1(a_1a_2 - a_3a_4) \\ 2a_3a_4^2 & -2a_4(a_2 - a_3)(a_2 + a_3) \\ -2a_1^2a_4 & -2a_1(a_1a_3 - a_2a_4) \\ -a_2a_4^2 & 2a_2(a_1a_2 - a_3a_4) \\ 2a_2^2a_4 & 2a_2(a_2a_3 - a_1a_4) \\ -2a_3a_4^2 & 2a_4(a_1 - a_3)(a_1 + a_3) \\ a_4^3 & -a_4(4a_1a_2 - 3a_3a_4) \\ 0 & 2a_4(a_1 - a_2)(a_1 + a_2) \\ a_4(4a_2a_3 - a_1a_4) & 2a_3(a_2a_3 - a_1a_4) \\ a_4(4a_1a_3 - a_2a_4) & 2a_3(a_1a_3 - a_2a_4) \\ -2a_1a_4^2 & -a_4(4a_1a_3 - 3a_2a_4) \\ -2a_2a_4^2 & -a_4(4a_2a_3 - 3a_1a_4) \end{bmatrix}$$

5.9. The Polynomials Defining the Line Scheme of $\mathcal{H}(\mathfrak{sl}_k(2, \mathbb{k}))$.

- 5.9.1. $-M_{14}(2M_{13}M_{12}^2 + M_{24}M_{12}^2 - M_{13}M_{14}^2 + M_{13}M_{14}M_{23} - M_{13}^2M_{24} + M_{14}M_{23}M_{24})$
- 5.9.2. $M_{12}^2M_{14}^2 - M_{23}^2M_{14}^2 + M_{13}M_{24}M_{14}^2 - M_{23}M_{24}^2M_{14} - M_{12}^2M_{24}^2 + M_{13}^2M_{24}^2$
- 5.9.3. $-M_{14}M_{24}(M_{13}M_{14} - M_{23}M_{24})$
- 5.9.4. $M_{24}(M_{14}M_{12}^2 + 2M_{23}M_{12}^2 - M_{14}M_{23}^2 - M_{23}M_{24}^2 + M_{13}M_{14}M_{24} + M_{13}M_{23}M_{24})$
- 5.9.5. $2M_{12}M_{14}M_{24}M_{34}$
- 5.9.6. $2M_{14}^2M_{24}M_{34}$
- 5.9.7. $2M_{14}M_{24}^2M_{34}$
- 5.9.8. $M_{14}M_{23} - M_{13}M_{24} + M_{12}M_{34}$
- 5.9.9. $-M_{12}M_{14}(2M_{12}M_{13} + M_{34}M_{13} + M_{12}M_{24})$
- 5.9.10. $M_{14}^2(2M_{12}M_{13} + M_{34}M_{13} + M_{12}M_{24})$
- 5.9.11. $M_{14}(2M_{12}M_{13}M_{23} - M_{12}M_{24}M_{23} - M_{13}M_{34}M_{23} - M_{12}M_{14}M_{24} + M_{13}M_{14}M_{34})$
- 5.9.12. $-M_{14}(M_{34}M_{12}^2 - 2M_{13}^2M_{12} + M_{14}^2M_{12} + M_{14}M_{23}M_{12} - M_{13}^2M_{34} + M_{14}M_{23}M_{34})$
- 5.9.13. $M_{14}(M_{34}M_{12}^2 + M_{14}^2M_{12} + M_{14}M_{23}M_{12} + M_{13}M_{24}M_{12} + M_{14}M_{23}M_{34})$
- 5.9.14. $-M_{14}M_{24}(2M_{12}M_{13} + M_{34}M_{13} + M_{12}M_{24})$
- 5.9.15. $M_{14}M_{24}(2M_{12}M_{13} - M_{34}M_{13} + M_{12}M_{24})$

- 5.9.16. $-M_{14}M_{34}M_{12}^2 - M_{23}M_{34}M_{12}^2 - M_{14}M_{23}^2M_{12} + M_{14}M_{24}^2M_{12} + M_{23}M_{24}^2M_{12}$
 $- M_{13}M_{23}M_{24}M_{12} + M_{14}M_{23}^2M_{34} - M_{13}M_{14}M_{24}M_{34}$
- 5.9.17. $M_{12}M_{24}(M_{12}M_{14} + 2M_{12}M_{23} - M_{23}M_{34})$
- 5.9.18. $M_{12}(M_{13}M_{14} - 2M_{13}M_{23} + M_{23}M_{24})M_{34}$
- 5.9.19. $-M_{24}(M_{12}M_{13}M_{14} + M_{12}M_{24}M_{14} - M_{13}M_{34}M_{14} - 2M_{12}M_{13}M_{23} + M_{13}M_{23}M_{34})$
- 5.9.20. $M_{13}(M_{13}M_{14} - 2M_{13}M_{23} + M_{23}M_{24})M_{34}$
- 5.9.21. $M_{14}M_{24}(M_{12}M_{14} + 2M_{12}M_{23} - M_{23}M_{34})$
- 5.9.22. $-M_{13}M_{34}M_{12}^2 - M_{24}M_{34}M_{12}^2 - M_{13}M_{14}^2M_{12} + M_{13}M_{14}M_{23}M_{12} + M_{13}^2M_{24}M_{12} -$
 $M_{14}^2M_{24}M_{12} + M_{13}^2M_{24}M_{34} - M_{14}M_{23}M_{24}M_{34}$
- 5.9.23. $M_{14}M_{24}(M_{12}M_{14} + 2M_{12}M_{23} + M_{23}M_{34})$
- 5.9.24. $-M_{14}(2M_{12}M_{13}M_{23} - M_{12}M_{24}M_{23} + M_{13}M_{34}M_{23} - M_{24}M_{34}M_{23} - M_{12}M_{14}M_{24})$
- 5.9.25. $M_{23}(M_{13}M_{14} - 2M_{13}M_{23} + M_{23}M_{24})M_{34}$
- 5.9.26. $M_{24}(M_{34}M_{12}^2 - M_{24}^2M_{12} - M_{14}M_{23}M_{12} - M_{13}M_{24}M_{12} + M_{13}M_{24}M_{34})$
- 5.9.27. $M_{24}(M_{34}M_{12}^2 + 2M_{23}^2M_{12} - M_{24}^2M_{12} - M_{13}M_{24}M_{12} - M_{23}^2M_{34} + M_{13}M_{24}M_{34})$
- 5.9.28. $M_{24}^2(M_{12}M_{14} + 2M_{12}M_{23} - M_{23}M_{34})$
- 5.9.29. $-M_{24}(M_{12}M_{13}M_{14} + M_{12}M_{24}M_{14} - 2M_{12}M_{13}M_{23} - M_{13}M_{23}M_{34} + M_{23}M_{24}M_{34})$
- 5.9.30. $-M_{14}M_{34}(2M_{12}M_{13} + M_{34}M_{13} - M_{12}M_{24})$
- 5.9.31. $M_{14}^2(M_{14}^2 - M_{24}^2 - M_{34}^2)$
- 5.9.32. $-M_{14}M_{24}M_{13}^2 + M_{23}M_{24}M_{13}^2 - M_{14}M_{23}^2M_{13} + M_{14}M_{34}^2M_{13} - M_{23}M_{34}^2M_{13}$
 $+ M_{12}M_{23}M_{34}M_{13} + M_{14}M_{23}^2M_{24} - M_{12}M_{14}M_{24}M_{34}$
- 5.9.33. $-M_{14}M_{34}(M_{12}M_{14} + M_{23}M_{34})$
- 5.9.34. $-M_{13}^2M_{14}^2 + M_{23}^2M_{14}^2 - M_{12}M_{34}M_{14}^2 - M_{23}M_{34}^2M_{14} - M_{12}^2M_{34}^2 + M_{13}^2M_{34}^2$
- 5.9.35. $-M_{24}M_{34}(M_{12}M_{24} - M_{13}M_{34})$
- 5.9.36. $-M_{13}^2M_{24}^2 + M_{23}^2M_{24}^2 - M_{12}M_{34}M_{24}^2 + M_{13}M_{34}^2M_{24} + M_{12}^2M_{34}^2 - M_{23}^2M_{34}^2$
- 5.9.37. $M_{14}M_{24}(M_{14}^2 + M_{23}M_{14} - M_{24}^2 - M_{34}^2 - M_{13}M_{24} - M_{12}M_{34})$
- 5.9.38. $M_{14}M_{24}(M_{14}^2 + M_{23}M_{14} - M_{24}^2 + M_{34}^2 - M_{13}M_{24} - M_{12}M_{34})$
- 5.9.39. $-M_{24}M_{34}(M_{12}M_{14} - 2M_{12}M_{23} + M_{23}M_{34})$
- 5.9.40. $-M_{14}M_{24}M_{13}^2 + M_{23}M_{24}M_{13}^2 - M_{14}M_{23}^2M_{13} + M_{23}M_{34}^2M_{13} + M_{12}M_{23}M_{34}M_{13} -$
 $M_{23}M_{24}M_{34}^2 + M_{14}M_{23}^2M_{24} - M_{12}M_{14}M_{24}M_{34}$
- 5.9.41. $M_{24}^2(M_{14}^2 - M_{24}^2 + M_{34}^2)$
- 5.9.42. $-M_{34}(M_{24}M_{13}^2 - M_{34}^2M_{13} + M_{14}M_{23}M_{13} - M_{12}M_{34}M_{13} + M_{12}M_{24}M_{34})$
- 5.9.43. $-M_{14}M_{34}(M_{14}^2 - M_{23}M_{14} + M_{24}^2 - M_{34}^2 - M_{13}M_{24} - M_{12}M_{34})$

- 5.9.44. $-M_{34}(M_{14}M_{23}^2 - M_{34}^2M_{23} + M_{13}M_{24}M_{23} + M_{12}M_{34}M_{23} - M_{12}M_{14}M_{34})$
 5.9.45. $-M_{24}M_{34}(M_{14}^2 - M_{23}M_{14} + M_{24}^2 - M_{34}^2 - M_{13}M_{24} + M_{12}M_{34})$
 5.9.46. $-M_{34}^2(M_{14}^2 + M_{24}^2 - M_{34}^2)$

Acknowledgments. The authors gratefully acknowledge support from the NSF under grants DMS-0900239 and DMS-1302050.

REFERENCES

- [1] M. ARTIN, J. TATE AND M. VAN DEN BERGH, Some Algebras Associated to Automorphisms of Elliptic Curves, in “The Grothendieck Festschrift 1”, pp 33-85, Eds. P. Cartier et al, Birkhäuser Boston (1990).
- [2] M. ARTIN, J. TATE AND M. VAN DEN BERGH, Modules Over Regular Algebras of Global Dimension 3, *Invent. Math.* **106** (1991), 333-388.
- [3] R. CHANDLER, On the Quantum Spaces of some Quadratic Regular Algebras of Global Dimension Four, *Ph.D. Thesis, University of Texas at Arlington* (2016).
- [4] R. CHANDLER AND M. VANCLIFF, The One-Dimensional Line Scheme of a Certain Family of Quantum \mathbb{P}^3 s, *J. Alg.* **439** (2015), 316-333.
- [5] X. W. CHEN, S. D. SILVESTROV, AND F. VAN OYSTAEYEN, Representations and Cocycle Twists of Color Lie Algebras, *Springer Sci.-Alg. Rep. Theory* **9** (2006), 633-650.
- [6] S. CHENG AND W. WANG, Dualities and Representations of Lie Superalgebras, Graduate Studies in Mathematics **144**. *Amer. Math. Soc., Rhode Island* (2013).
- [7] D. A. COX, J. B. LITTLE, AND D. O’SHEA, Ideals, Varieties and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra, *Springer-Verlag, New York* (1992).
- [8] L. D. FADDEEV, N. Y. RESHETIKHIN, AND L. A. TAKHTADZHIAN, Quantization of Lie Groups and Lie Algebras, *Leningrad Math. J.* **1** (1990), 193-225.
- [9] K. R. GOODEARL AND R. B. WARFIELD, JR., An Introduction to Noncommutative Noetherian Rings, Student Texts **61**, *Lond. Math. Soc.* (2004).
- [10] J. HARRIS, Algebraic Geometry: A First Course, Graduate Texts in Mathematics **133**, *Springer-Verlag, New York* (1992).
- [11] J. C. JANTZEN, Lectures on Quantum Groups, Graduate Studies in Math **6**. *Amer. Math. Soc., Rhode Island* (1996).
- [12] L. LE BRUYN AND S. P. SMITH, Homogenized $\mathfrak{sl}(2)$, *Proceedings of the Amer. Math. Soc.* **118** No. 3 (1993), 725-730.

- [13] L. LE BRUYN, S. P. SMITH AND M. VAN DEN BERGH, Central Extensions of Three-Dimensional Artin-Schelter Regular Algebras, *Math. Z.* **222** (1996), 171-212.
- [14] T. LEVASSEUR, Some Properties of Noncommutative Regular Graded Rings, *Glasgow Math. J.* **34** (1992), 277-300.
- [15] T. LEVASSEUR AND T. STAFFORD, The Quantum Coordinate Ring of the Special Linear Group, *J. Pure and Appl. Alg.* **86** (1993), 181-186.
- [16] B. SHELTON AND M. VANCLIFF, Schemes of Line Modules I, *J. Lond. Math. Soc.* **65** No. 3 (2002), 575-590.
- [17] B. SHELTON AND M. VANCLIFF, Schemes of Line Modules II, *Comm. Alg.* **30** No. 5 (2002), 2535-2552.
- [18] S. P. SMITH, Quantum Groups: An Introduction and Survey for Ring Theorists, in "Noncommutative Rings," *MSRI Publications* **24** (1992), 131-178.
- [19] M. VANCLIFF, Quadratic Algebras Associated with the Union of a Quadric and a Line in \mathbb{P}^3 , *J. Alg.* **165** (1994), 63-90.